



INCOBEST 2023

Universitas Muhammadiyah Surakarta
International Conference on Biology Education, Natural Science, and Technology

"Emerging Issues on Natural Science, Environment, and Its Learning Innovation"

PROCEEDINGS BOOK

International Conference on Biology Education, Natural Science, and Technology (INCOBEST)

June 17th, 2023 - Surakarta

DEPARTMENT OF BIOLOGY EDUCATION
FACULTY OF TEACHER TRAINING AND EDUCATION
UNIVERSITAS MUHAMMADIYAH SURAKARTA

incobest.ums.ac.id



Editor:

- **Vina Listiawati**, (SCOPUS ID: [57225089904](#)) University of the Ryukyus, Japan
- **Jumadi**, (SCOPUS ID: [57200103386](#)) Universitas Muhammadiyah Surakarta, Indonesia
- **Nazmul Hasan**, (SCOPUS ID: [57222598153](#)) University of Kentucky, USA
- **M. Luthfi Hidayat**, (SCOPUS ID: [58582981100](#)) King Abdulaziz University, Saudi Arabia

Reviewer:

- **Suranto**, (SCOPUS ID: [57208564292](#)) Universitas Muhammadiyah Surakarta, Indonesia
- **Martin Lamb**, (SCOPUS ID: [35958438300](#)) University of Leeds, United Kingdom
- **Kongkiti Phusavat**, (SCOPUS ID: [13403740100](#)) Kasetsart University, Thailand
- **Rita Pramujianti Khotimah**, (SCOPUS ID: [57210910953](#)) Universiti Pendidikan Sultan Idris, Malaysia
- **Mohamad Waluyo**, (SCOPUS ID: [57213869096](#)) Universitas Muhammadiyah Surakarta, Indonesia
- **Mariani Binti Md Nor**, (Scopus ID: [57190401743](#)) Universiti of Malaya, Malaysia
- **Michio Matsubara**, (Scopus ID: [7202863298](#)) Kanazawa University, Japan
- **Muhammad Kristiawan**, (Scopus ID: [57205367909](#)) Universitas Bengkulu, Indonesia
- **Tommy Tanu Wijaya**, (Scopus ID: [57218281226](#)) Guangxi Normal University, China
- **Sabariah Binti Sharif**, (Scopus ID: [39862198600](#)) University Malaysia Sabah, Malaysia

Keynote Speakers:

- **Assoc. Prof. Haniza Hanim**, (SCOPUS ID: [57208674877](#)) Universiti Pendidikan Sultan Idris, Malaysia
- **Prof. Luiz Gustavo Galego**, (SCOPUS ID: [12753764000](#)) Universidade Federal do Triangulo Mineiro, Brazil
- **Dr. Triastuti Rahayu, M.Si.**, (SCOPUS ID: [57384759900](#)) Universitas Muhammadiyah Surakarta, Indonesia

Steering Committee (Proceedings):

- **Yasir Sidiq**, (SCOPUS ID: [57215926493](#)) Universitas Muhammadiyah Surakarta, Indonesia
- **Guntur Nurcahyanto**, (Google Scholar ID: [Gh1aiAQAAAAJ](#)) Universitas Muhammadiyah Surakarta, Indonesia
- **Santhyami**, (SCOPUS ID: [57200796303](#)) Universitas Muhammadiyah Surakarta, Indonesia
- **Ambarwati**, (SCOPUS ID: [57210644042](#)) Universitas Muhammadiyah Surakarta, Indonesia
- **Efri Roziaty**, (SCOPUS ID: [57211964450](#)) Universitas Muhammadiyah Surakarta, Indonesia
- **Paramitha Cahyani**, (SCOPUS ID: [57424029100](#)) Universitas Muhammadiyah Surakarta, Indonesia

The Ecology of Zaprionus Genus in Brazil and Adaptation Process during The Bioinvasion

Prof. Luis Gustavo da Conceicao Galego

1-21

 PDF

Emerging Innovations in Natural Science Education

Haniza Hanim binti Mohd Zain, Ph.D.

22

 PDF

Metagenomic Studies and Their Applications

Dr. Triastuti Rahayu

23-32

 PDF

Development of Android-Based Interactive Teaching Materials Using the Ispring Application with Insight into Sustainable Development

N Nurhasnah, L Lufri, Abdul Razak, F Festiyed, Prima Aswirna, Windy Kasmita, Wetri Yesmoneca

33-41

 PDF

The Effect of STEM Approach on Students' Critical Thinking Skills

Media Roza, Minda Azhar, F Festiyed, Prima Aswirna, Rika Purnama Sari

42-51

 PDF

Inventory of Montane Zone Weeds in the Selo Tourism Area, Boyolali Regency, Central Java

Nila Permatasari, S Santhyami

52-61

 PDF

Development of Problem-Based HOTS (Higher Order Thinking Skill) Question Instruments on Class VIII Pressure Material

Rina Irawati, S Sukarmin, Sri Yamtinah

62-67

 PDF

The Characteristics of Herbal Tea Combination Between Butterfly Flower with Mint and Pandan Leaf on Drying Duration Variation

Titik Suryani, Ria Anisa Rahmasari, Riza Ainun Nisa

68-76

 PDF

Preservice Biology Teacher's TPACK Integration into Lesson Planning

Putri Agustina, Lina Agustina, Rina Astuti, Yesi Chotimah, Putri Gayatri Ramadhanty, Alanindra Saputra

77-83

 PDF

Students' Creative Thingking Ability in view From The Level of Understanding in Waste Management

Syarifah Zahrah, Rita Retnowati, Rita Istiana

84-90

 PDF

Monitoring The Correlation of Climatics to The Airborne Bacteria at The Manggarai Station, South Jakarta, Indonesia

Reza Anindita, Auliya Rahmawati, Maulin Inggraini, Melania Perwitasari, Dede Dwi Nathalia, Maya Uzia Beandrade, Intan Kurnia Putri

91-100

 PDF

The Effect of Active Learning Integrated with Everyone Is A Teacher Here Strategy Based on Questioning and Answering on The Critical Thinking Skills and Biology Learning Retention of Class XI Senior High School Students In The Industrial Agriculture Area

Kurnia Suciningsih, Bea Hana Siswati, Nadyatul Ilma Indah Savira

101-108

 PDF

The Effect of STEM (Science, Technology, Engineering, and Mathematics) based Textbooks in Biotechnology Learning Material on The Critical Thinking Skills and Learning Results of Senior High School Students in The Industrial Agriculture Area

Kamilatul Khoiroh, Bea Hana Siswati, Kamalia Fikri

109-121

 PDF

Quality of Moringa oleifera Leaf and Kiambang (Salvinia molesta) Solid Organic Fertilizer with Banana Peel Bioactivator

Desvia Puput Apriani, Aminah Asngad

122-132

 PDF

Quality of Solid Organic Fertilizers for Duck Manure and Trembesi Leaves with Papaya Peel Bioactivator

Inventory of Moss Plants (Bryophyta) in the Montana Zone, Samiran Village, Selo District, Boyolali Regency, Central Java Province

Eva Diahayu Wardani, S Santhyami

143-151

 PDF

Diversity of Ferns (Pteridophyta) in the Montana Zone, Samiran Village, Selo District, Boyolali Regency, Central Java Province

Indah Rahmawati, S Santhyami

152-159

 PDF

Quality of Solid Organic Fertilizer from Moringa Leaves and Peanut Shells with Banana Peel Bioactivator

Nur Afifah Mustikasari, Aminah Asngad

160-170

 PDF

Quality of Liquid Organic Fertilizer Peanut Shells and Tofu Dregs With Banana Stems as Bioactivator

Muhammad Rizal Kahfi, Aminah Asngad

171-180

 PDF

The Quality of Infrastructure Facilities and the Readiness of Biology Laboratories to Support Learning in High School

Mei Renni Sholikhah, Lina Agustina

181-192

 PDF

Quality of Duck Manure Fertilizer and Paitan Leaves with the Addition of MOL Papaya Skin

Hana Khairunnisa, Aminah Asngad

193-203

 PDF

Quality of Liquid Organic Fertilizer Combination Gamal Leaves and Tofu Dregs with Banana Stem Bioactivator

Feby Istifarini, Aminah Asngad

204-214

 PDF

The Effects of Socio-Scientific Approach on Junior High School Students' Learning Outcomes

Salsabila Nur Rahmadhani, Putri Agustina, Sri Mulyani

215-226

 PDF

Study of Science Practicum in Junior High School, Sukoharjo, Indonesia

Farikha Muya Sarah, Putri Agustina, Harini Hastowati

227-240

 PDF

Quality of Kefir Combination Between Soy Milk and Skim Milk On Variation of Sugar and Fermentation Duration

Masithoh Uswatiningtyas, Titik Suryani

241-247

 PDF

Coconut Husks and Eggshells Liquid Organic Fertilizer's Effect on The Growth and Calcium Levels of Kale

Nabila Junisky Susetyawati, Aminah Asngad

248-259

 PDF

Phosphate Solubilisation Index and Antagonism Potential of Frangipani Tree Rhizosphere Bacterial Isolates from Cemetery

Vika Febriyanti, Erma Musbita Tyastuti, Yasir Sidiq, Triastuti Rahayu

260-268

 PDF

Coconut Husk And Eggshell Liquid Organic Fertilizer Effect On Lettuce Growth And Calcium Levels

Widya Husnul Khotimah, Aminah Asngad

269-279

 PDF

Quality Of Kirinyuh Leaf Liquid Organic Fertilizer and Coconut Water Waste With Pineapple Skin Bioactivator

Arisa Putri Jihan Nitami, Aminah Asngad

280-290

 PDF

Isolation and Characterization of Cellulolytic Bacteria from Bonoloyo, Cemetery

Wanda Datik Risnasari, Erma Musbita Tyastuti, Yasir Sidiq, Triastuti Rahayu

291-298

 PDF

Application of Project Based Learning Model to Improve Critical Thinking Skills of Students SMA Muhammadiyah 1 Sragen

Tasya Cahya Dwi Purwanti, Rina Astuti

299-304

 PDF

Quality of Kefir Combination Between of Goat's Milk and Skim Milk on Variations of Sugar and Fermentation Duration

Erna Kurniawati, Titik Suryani

305-313

 PDF

Quality of Herbal Tea Combination Between Corn Silk and Stevia Leaf on Variation of Temperature and Drying Duration

Feny Nafiza, Titik Suryani

314-325

 PDF

Isolation and Identification of Soil Bacteria in Pracimaloyo Public Cemetery, Kartasura

Bagas Adityaradja, Erma Musbita Tyastuti, Yasir Sidiq, Triastuti Rahayu

326-334

 PDF

LOF Kirinyuh Leaves and Liquid Waste Tempeh with Pineapple Peel Bioactivator: Sensory and Macronutrient Content

Nur Jati Sekaringsih, Aminah Asngad

335-347

Glucose Levels and Organoleptic Quality Probiotic Tepache of Pineapple Peel on Variation of Sugar and Fermentation Duration
Qurota Ayuni Nuha Sabira, Titik Suryani 348-355

 PDF

Empowerment of Animalia Classification Material: A Review of Literature
Tunjung Nala Puti, Puguh Karyanto, Umi Fatmawati 356-367

 PDF

Quality of Herbal Tea Combination Between Corn Silk and Mint Leaf in a Variation of Temperature and Drying Duration
Imroatul Husna, Titik Suryani 368-374

 PDF

Number of Erythrocytes of White Rats (*Rattus norvegicus*) Sonded with a Combination of Porang Tuber and Moringa Leaf Extract
Alya Anggitasari, Endang Setyaningsih 375-384

 PDF

Pakcoy (*Brassica chinensis* L.) Growth Using Moringa Leaf Extract with the Addition of Red Onion Skin
Dwi Putri Widjaya, S Suparti 385-393

 PDF

Mustard (*Brassica Juncea* L.) Growth Hydroponically Using AB-Mix and Liquid Organic Fertilizer Tea Pulp
Dinda Putri Trisnawati, S Suparti 394-401

 PDF

The Effect of Cassava Peel Powder on the Productivity of the Brown Oyster Mushroom (*Pleurotus cystidiosus*)
Emmalia Rahmawati, S Suparti 402-410

 PDF

Analysis of Misconceptions on Reproductive System Material in Class XI IPS 2 Students of SMA Negeri 6 Madiun
Jevi Milda Rahmawati, Ima Agung Nurcahyo, Cicilia Novi Primiani 411-420

 PDF

Inventory of Epiphytic Lichen Foliose on Tea Plants in Ngargoyoso, Karanganyar, Central Java
Dika Dwi Irnawati, Efri Rozlaty 421-429

 PDF

Manuscript Writing Guidelines: Analysis of Creative thinking skills profile of Senior High school students in Madiun City on Biotechnology topic
Herning Nurdiana, Ima Agung Nurcahyo, Cicilia Novi Primiani 430-437

 PDF

Plasma Protein of Obese White Rats Treated with of Glucomannan Porang Tubers and Moringa Extract
Sabila Dina Ikhsani, Endang Setyaningsih 438-445

 PDF

Effectiveness of Flipped Classroom Model with Instagram in Junior High School
Mega Amartya Istighfarin, Lina Agustina 446-454

 PDF

Application of the NHT Model Combined with Question Cards on Biology Learning Outcomes

Nabilla Riska Nurjannah, D Djumadi

455-463

 PDF

Growth Red Spinach (*Amaranthus amoenus*) by Hydroponics Using Charcoal Media Husk

I Isherdini, Dyah Salma Nurlaila, S Suparti

464-472

 PDF

Oxygen Consumption and BB *Rattus norvegicus* Obesity Sonde Treatment Glucomanan Porang Tubers and Moringa Leaf Extract

Mustika Arum Mayang Sari, Endang Setyaningsih

473-483

 PDF

The Effect of Kepok Banana Skin Liquid Organic Fertilizer (*Musa paradisiaca*) on the Growth of Mustard Plants (*Brassica juncea* L.)

I Isherdini, Berliana Putri Athena, S Suparti

484-488

 PDF

The Effect of PBL Learning Model Combined with Rewards and Punishment on Science Learning Outcomes

Annisa Tri Yudhatami, D Djumadi

489-501

 PDF

Antioxidant Activity and Organoleptic Quality of Probiotic Tepache of Pineapple Peel Sugar Variation and Fermentation Duration

Fithrotul Haq Rabbani Abdurrahman, Titik Suryani

502-508

 PDF

Analysis of Plant Vegetation in Sekipan Forest Tawangmangu District, Karanganyar District

Ima Aryani, Shal Syabela Riskawati

509-517

 PDF

Health Examination of The Digestion of Proboscis Monkey (*Nasalis larvatus*) through Bacteriological Tests on Feces

Camalia Maisya, Anni Nurliani, Amalia Rezeki, R Rusmiati

518-532

 PDF

The Phylogeny of Parasitic Hemiptera Using Multivariate Statistical Package Cluster Analysis

Dwi Setyo Astuti, Murwanto Setyo Nugroho

533-540

 PDF

Identification of Intestine Parasite Worms Eggs in Feces Proboscis Monkey (*Nasalis larvatus*)

Fiqita Giulleta Irtifannisa Rahman, Anni Nurliani, Amalia Rezeki, R Rusmiati

541-549

 PDF

Ethnobotanical Study of Tea Plants Based on Local Wisdom in Ngargoyoso District, Karanganyar Regency, Central Java

Kharisma Margiyana Dinda Hapsari, Efri Roziaty

550-556

 PDF

Scientific Learning with PjBL and PBL Models of Science Process Skills Student Learning Outcomes

Farid Nur Yusron, S Sarwanto, Mohammad Masykuri

557-563

 PDF

Water Quality in Satui River and Histological Structure Study of Liver of Baung Fish (*Mystus nemurus*)

Dinda Triana, Heri Budi Santoso, Anang Kadarsah

564-571

 PDF

Profile of Creative Thinking Skill of New Normal Era at Junior High School Students

Farida Amrul Almuharomah, Widha Sunarno, Mohammad Masykuri

572-579

 PDF

The Feasibility and Effectiveness of High-Level Thinking Tests on Virus to Mapping The Ability of Prospective Science Teacher (Introduction Study on The Development of Integrated Learning Models)

Annur Indra Kusumadani, Sentot Budi Rahardjo, Sri Yamtinah, Baskoro Adi Prayitno

580-586

 PDF

PBL-Based Science Learning on Living Things Interaction in Junior High School

H Hariyatmi, Hidayatus Sholihah Tisniasari

587-598

 PDF

Monitoring The Correlation of Climatics to The Airborne Bacteria at The Manggarai Station, South Jakarta, Indonesia

Reza Anindita^{1*}, Auliya Rahmawati², Maulin Inggraini³, Melania Perwitasari⁴, Dede Dwi Nathalia⁵, Maya Uzia Beandrade⁶, Intan Kurnia Putri⁷

^{1,4,5,6,7}Department of Pharmacy, Stikes Mitra Keluarga, East Bekasi, Indonesia

^{2,3}Department of Medical Laboratory Technology, STIKes Mitra Keluarga, East Bekasi, Indonesia

*Corresponding Author. E-mail address: rezaanindita@gmail.com

ABSTRACT

KEYWORDS:

Temperature,
humidity,
light intensity,
manggarai station,
bacteria

The highest of microorganisms suspended in the air were bacteria with a percentage of 80.8%. If the intensity of bacterial exposure occurs in large quantities, it will accumulate in the respiratory tract which has the potential to trigger allergic reactions and respiratory infections. Given the problems and impacts caused by air microorganisms on public health, it is necessary to monitor the distribution of air bacteria. The purpose of this study was to determine the relationship between temperature and humidity with the number of bacteria in the Manggarai station area. Type of research is correlation. The sampling location is Manggarai Station, East Jakarta, Indonesia. The 10 sampling points in this study include the station hall, motorbike parking, prayer rooms, train platform 1-2, train platform 2-3, train platform 3-4, and train platform 5, toilet, and underpass. The results of this study included the highest number of bacteria found in the station hall 331 colonies, the highest percentage of bacterial morphology is monobacilli was 78%, the distribution of gram positive was more than gram negative with spore bacteria being more dominant than non spore. The correlation coefficient between temperature and number of bacterial colonies (0.134) and humidity with number of bacterial colonies (0.380) showed weak positive correlation, while the correlation coefficient for light intensity with the number of bacterial colonies - 0.140 (very weak negative).

© 2023 The Author(s). Published by Biology Education Department, Faculty of Teacher Training and Education, Universitas Muhammadiyah Surakarta. This is an open access article under the CC BY-NC license: <https://creativecommons.org/licenses/by-nc/4.0/>.

1. INTRODUCTION

According to Bragoszewska and Pastuszka (2018), the percentage of airborne microorganisms is 10% of the 24% of the total particles suspended in the air. Cho *et al.* (2019) reported that the highest types of microorganisms suspended in the air were bacteria with a percentage of 80.8%. The dominant bacterial phyla identified in the air were Proteobacteria (32.2%), Cyanobacteria (18.0%), Actinobacteria (16.5%), Firmicutes (15.5%), and Bacteroidetes (11.6%) (Liu *et al.*, 2018).

Kallawicha *et al.* (2015) explained that bacteria are components of bioaerosols that are abundant in the air both outdoors and indoors. In addition to bacteria, other bioaerosol components are viruses, fungi, pollen, metabolites of microorganisms (mycotoxins), and endotoxins as the outer membrane of bacterial cells that are released during bacterial lysis and growth. The presence of bacteria and endotoxins as components of bioaerosols were identified as important factors affecting human health. According to Jones and Harrison (2004) bacteria have a size of 0.25–8 m. This size makes it easier for bacteria to enter the human body through inhalation of the respiratory tract. If the intensity of bacterial exposure occurs in large quantities, it will accumulate in the respiratory tract which has the potential to trigger allergic reactions and respiratory infections such as asthma, rhinitis, pneumonia, and atopic dermatitis.

Cho *et al.* (2019) reported that bacteria with a size of 5-10 μ m that accumulate in the upper respiratory tract can trigger rhinitis, while bacteria with a size of $<5 \mu$ m that accumulate in the alveoli can trigger allergic reactions. However, these reactions vary between individuals. This effect tends to be more dangerous in someone with a weakened, moderate, or weakened immune system, such as children, pregnant women, and the elderly.

Given the problems and impacts caused by air microorganisms on public health, it is necessary to monitor the distribution of air microorganisms. Monitoring can be done by conducting research on the relationship between physical quality and outdoor air bacteria, especially at stations as one of the centers of crowds in densely populated urban areas.

This research was conducted at the Manggarai Station in South Jakarta, Indonesia. This station is a transit center for trains that is full of people and is a source of spreading airborne bacteria in the outside environment through coughing, breathing, and exposure to human skin. The selection of research that focuses on outdoor air bacteria at the Manggarai station refers to previous studies which were more dominant in monitoring indoor air bacteria.

Several previous studies that examined the relationship between temperature and humidity on air microorganisms, among others, research Menteşe *et al.* (2009); Wu *et al.* (2012); Kumar *et al.* (2011); Bragoszewska and Pastuszka (2018); Cho *et al.* (2019); Goudarzi *et al.*, (2017). The difference between this research and previous research is in terms of location, this research was conducted in Indonesia, especially at the Manggarai Station. The novelty of this research is that there has never been data on air bacteria in the outdoor environment in the Manggarai station area, East Jakarta, Indonesia, considering that research on air bacteria in Indonesia is more dominant indoors.

The purpose of this study was to determine the relationship between temperature and humidity with the number of bacteria in the Manggarai station area. The existence of data on airborne bacteria can be used as a reference in formulating better policies regarding Indonesian public health guidelines outdoors.

2. MATERIALS AND METHODS

This type of research is a correlation. The sampling location is Manggarai Station, East Jakarta, Indonesia. Samples were examined at the Microbiology Laboratory of STIKes Mitra Keluarga, East Bekasi.




Figure 1. Map manggarai station location, south jakarta, indonesia

The instrument used in this study include Autoclave (Hirayama HG-50), Incubator (DNP), Analytical Balance (Adam), Microscope (Olympus CX22), Showcase (polytron), BSC (JSR), Hot plate and stirrer (Ika HS-10), Colony counter (KJY-020), petri dish (pyrex), bunsen's burner, Erlenmeyer (pyrex), spatula, beaker (pyrex), Thermohygrometer, Lux meter, tripod, needle loop, stirring rod, matches, dropper, object glass (sail brand) and coloring tub. The materials used in this study were 70% alcohol, spirits, aquadest, Nutrient Agar media (Himedia), Crystal violet (Be Reagent), Lugol's iodine (Merck), 95% alcohol, safranin, and Malachite Green (Riedel-De Haen).

The population in this study is the entire area of the Manggarai station. Determination of the sample area in this study using a purposive sampling technique with the criteria of a crowded area and frequently visited by passengers. The 10 sampling points in this study include the station hall, motorbike parking, prayer rooms, train platform 1-2, train platform 2-3, train platform 3-4, and train platform 5, toilet, and underpass.

The procedure of this research include pre-analytic, analytic, and post-analytic stages :

The Pre-analytic

The pre-analytic stage starts by making Nutrient Agar (NA) media to grow bacteria, measuring temperature and humidity with a thermohygrometer, light intensity with a lux meter, and taking bacterial samples at 10 points in the Manggarai station area. Each point is repeated three times on the same day and at different times, at 07.00 - 09.00 WIB. This time is the departure time of passengers at Manggarai station based on the results of a preliminary survey. The air sampling process is carried out using NA media that has been coded in the form of date, time, and location of collection placed in a predetermined sampling area with a height of 1.5 m. The NA medium was placed in an open petri dish for 10-15 minutes. The NA medium was closed again and incubated at 37°C for 2x24 hours. After incubation, the NA medium was observed and the number of bacterial colonies counted using colony counter.

The Analytic

The analytical stage is laboratory tests which include gram staining and spores. The staining results were observed using a light microscope with a magnification of 100x.

The post-analytic

The post-analytic stage is in the form of recording data on temperature, humidity, light intensity, number of bacteria, and types of bacteria based on morphology, gram, and presence of spores. All data were then analyzed by the Spearman Correlation test to determine the relationship between air physical factors such as temperature, humidity, and light intensity with the number of bacterial colonies.

The research has been conducted an ethical feasibility test by the Ethical Committee for Research health of Jakarta III Ministry of Health Polytechnic (KEPK-PKKJ3) with certificate ethic No. KEPK-PKKJ3 / 5 /II /2019

3. RESULTS AND DISCUSSION

3.1. Results

The results of examination bacteria after 2x24 hours incubation in this study showed that all areas showed the growth of bacterial colonies on petri dishes containing NA media. The results of the growth of bacterial colonies can be seen in **Figure 2**.




Figure 2. Bacterial colonies on NA media placed at the Manggarai station

The results of the calculation of the number of bacterial colonies after incubation of 2x24 NA media placed in 10 areas of Manggarai station in this study can be seen in **table 1**.

Table 1. The results of the calculation of the number of bacterial colonies

No	Areas	Colonies (CFU's)
1	The station hall	331
2	motorbike parking	189
3	Woman prayer rooms	33
4	Man prayer rooms	11
5	train platform 1-2	224
6	train platform 2-3	128
7	train platform 3-4	134
8	train platform 5	220
9	Woman toilet	169
10	Underpass	210




Figure 3. The number of bacterial colonies at Manggarai station

Table 1. shows that the average interval of the number of bacterial colonies found in 10 areas at the Manggarai station is between 11-331 colonies. The lowest number of bacterial colonies were found in the man prayer room as many as 11 colonies, while the highest was found in the hall area as many as 331 colonies (**Figure 3**). The distribution percentage of the number of bacteria based on their morphology can be seen in **Table 2** and **Figure 4**.

Table 2. The distribution percentage of the number of bacteria based on their morphology

Areas	Percentage of the number bacteria (%)								
	<i>Monobacilli</i>	<i>Diplobacilli</i>	<i>Streptobacilli</i>	<i>Coccobacilli</i>	<i>Monococci</i>	<i>Diplococci</i>	<i>Streptococci</i>	<i>Staphylococci</i>	<i>Sarcina</i>
The station hall	14	2	6	0	4	2	0	4	0
motorbike parking	12	7	7	1	8	5	10	10	1
Woman prayer rooms	3	0	0	0	1	0	0	0	0
Man prayer rooms	6	8	8	2	2	0	0	7	0
train platform 1-2	10	3	3	1	2	1	1	7	1
train platform 2-3	1	2	2	4	2	1	0	2	0
train platform 3-4	4	1	1	0	8	0	1	7	0
train platform 5	9	1	1	0	6	1	4	7	0
Woman toilet	7	1	1	0	7	1	2	4	0
Underpass	12	2	2	0	6	2	0	12	0
Total	78%	27%	31%	8%	46%	13%	18%	60%	2%


**Figure 4.** Diagram of the percentage number of bacterial colonies based on morphology

Table 2. show that there are 9 bacterial morphologies found in 10 areas of the Manggarai station, namely monobacilli, diplobacilli, streptobacilli, coccobacilli, monococci, diplococci, streptococci, staphylococci, and sarcina. The percentage of bacterial morphology that had the highest number was monobacilli, while the lowest was sarcina (**Figure 4**).

The results of the distribution of the number of bacteria based on gram staining can be seen in **Table 3.** and **Figure 5.**

Table 3. Distribution of bacteria based on gram staining

Area	Gram	
	Positive	Negative
The station hall	12	6
motorbike parking	19	7
Woman prayer rooms	2	2
Man prayer rooms	9	5
train platform 1-2	12	6
train platform 2-3	11	2
train platform 3-4	13	6
train platform 5	17	9
Woman toilet	1	15
Underpass	1	22




Figure 5. Distribution of bacteria based on gram staining




Figure 6. Gram-positive (A) and Gram-negative (B) bacteria.

In this study, the number of bacterial colonies based on the presence/absence of spores showed that the number of bacteria that did not have spores was more dominant than bacteria that had spores. The results of the number of bacterial colonies based on the presence or absence of spores are shown in **table 4** and **figure 7**.

Table 4. The number of bacterial colonies based on spores.

Area	Spore	
	Spore	Non-spore
The station hall	12	6
motorbike parking	12	14
Woman prayer rooms	3	1
Man prayer rooms	6	8
train platform 1-2	9	9
train platform 2-3	5	8
train platform 3-4	3	16
train platform 5	3	23
Woman toilet	5	11
Underpass	11	12




Figure 7. Diagram of the distribution of bacteria based on the presence of spores

The results in **table 4** and **figure 8** show that the percentage of bacteria that have spores in the hall (67%), motorbike parking (46%), women's prayer room (75%), men's prayer room (43%), platform 1-2 (50%), platform 2-3 (38%), platform 3-4 (16%), platform 5 (12%), women's toilet (31%), and underpass (48%), while the percentage of bacteria that did not have spores in the hall (33%), motorbike parking (54%), women's prayer room (25%), men's prayer room (57%), platform 1-2 (50%), platform 2-3 (62%), platform 3-4 (84%), platform 5 (88%), women toilets (69%), and underpass (52%). The microscopic picture of the structure of spore and non-spore bacteria can be seen in **Figure 8**.




Figure 8. Spore bacteria (a) and bakteri non-spore bacteria (b) (1000x).

In this study, measurements of physical factors at Manggarai Station were carried out simultaneously with air sampling at Manggarai Station. Measurement of physical factors was carried out at 10 sampling points at Manggarai Station. The data obtained consisted of temperature, humidity, and light intensity which can be seen in **table 5**.

Table 5. Results of measuring physical factors

Area	Temperature (°C)	Humidity (%)	Light Intensity (lux)
The station hall	31.8	63	405
motorbike parking	32.4	61	9324
Woman prayer rooms	30.1	64	1015
Man prayer rooms	29.8	66	83
train platform 1-2	31.6	63	3613
train platform 2-3	31.7	62	4100
train platform 3-4	32.1	60	2865
train platform 5	31.8	61	3057
Woman toilet	31	63	260
Underpass	31.1	63	67

Table 5. shows that the interval of the average temperature is between 29.8°C-32.4°C, humidity 60%-66%, and light intensity 83-9324 lux. The results of statistical tests on the correlation of physical factors of air quality toward the number of bacterial colonies using the Spearman correlation test with a 95% significance level can be seen in **table 6**.

Table 6. Correlation test results of temperature, humidity, and light intensity on the number of bacterial colonies

Test variable	Result	correlation coefficient
Temperature with the number of bacterial colonies	p = 0,000	r = 0,134
Humidity with the number of bacterial colonies	p = 0,000	r = 0,380
Light Intensity with the number of bacterial colonies	p = 0,000	r = -0,141

Table 6. shows that all test variables produce a p-value <0.05 or that there is a correlation between physical factors of air quality and the number of bacterial colonies. The correlation coefficient value between temperature and the number of bacterial colonies and humidity with the number of bacterial colonies showed a weak positive correlation, respectively 0.134 and 0.380, while the correlation coefficient (r) for light intensity with the number of bacterial colonies was -0.140 or classified as correlation. very weak negative.

3.2. Discussion

The results of the Spearman correlation test in this study showed that there was a correlation between temperature and the number of bacterial colonies at the Manggarai station ($P < 0.05$). However, this correlation is included in the very weak category ($r = 0.134$). The results of this study are in accordance with the research of Bragoszewska and Pastuszka (2018); Saadati *et al.* (2022) reported that there was a positive correlation between air temperature and the number of bacterial colonies.

According to Zhong *et al.* (2016), the correlation between temperature has a weak positive correlation with the number of bacterial colonies due to the presence of too high a temperature which can also inhibit bacterial growth. This is shown in the research results of Wu *et al.* (2012) who reported that increasing the temperature of UV light can significantly reduce the number of bacterial colonies outdoors. Therefore, the correlation between temperature and the number of bacterial colonies is very weak, considering that each bacterium has varying optimum temperature characteristics.

Another factor that correlates with the number of bacterial colonies is humidity. The correlation between humidity and the number of bacterial colonies in this study resulted in a positive correlation with a weak category. The results of this study are in accordance with the research of Kallawicha *et al.* (2015); Hwang and Yoon (2017) who reported that air humidity was positively correlated with the number of bacterial colonies. The difference in previous studies is that the correlation is strong, while in this study it is weak.

Hwang and Yoon (2017); Zhong *et al.* (2016); Hiwar *et al.* (2021) stated that humidity is related to the amount of water vapor in the air needed to increase cell wall strength and bacterial metabolism so that increasing air humidity can increase bacterial growth. However, humidity that is too high is often accompanied by very low temperatures that can inactivate protein enzymes that play a role in bacterial growth. Therefore, the correlation of humidity with the number of bacterial colonies often results in a weak positive correlation.

In contrast to temperature and humidity, in this study, the correlation between light intensity and the number of bacterial colonies resulted in a very weak negative correlation. The results of this study are in accordance with the research of Kamel *et al.* (2016); Fithri *et al.* (2016) who reported that light intensity was weakly negatively correlated with bacterial colonies. This is because some bacteria can have mutations that make them resistant to light exposure.

Gola *et al.* (2019) added that light intensity has bactericidal activity and plays an important role in spontaneous sterilization in natural conditions because sunlight contains ultraviolet light. Examples of bacteria that die from exposure to high light intensity are *Streptococcus* and other bacteria that cause respiratory tract infections.

Research result Gladka *et al.* (2021) ; Santos *et al.* (2011) ; Mohana *et al.* (2013) explained that although light intensity can inhibit the growth of some airborne bacteria, there are some bacteria that are resistant to high light intensity, such as *Micrococcus* sp. The resistance of these bacteria is obtained because some bacteria are able to form spores to survive and grow to spread to the environment without being affected by the intensity of sunlight. Therefore, the negative correlation between light intensity and the number of bacterial colonies is weak.

In this study, areas with a high number of bacterial colonies were station halls, platforms, and underpasses. The high number of bacteria is due to the fact that these three areas are the center of crowds at the station, thereby increasing the number of bacterial colonies carried by dust and humans as station users (Fang *et al.*, 2007); (Maier *et al.*, 2010).

The results of this study also showed that the dominant forms of airborne bacteria in the Manggarai station area were coccus and bacillus. These results are in accordance with the research of Menteşe *et al.* (2009) which states that the most common bacteria found in outdoor areas are coccus and bacillus with the dominant genus *Micrococcus* sp. *Streptococcus* sp and *Bacillus* sp. Goudarzi *et al.* (2017) added that in addition to the dominant fungal genus bacteria in the outdoor, among others, *Aspergillus* sp., *Penicillium* sp., and *Cladosporium* sp.

The results of this study also found the spread of the number of gram-positive and spore-forming bacterial colonies in large numbers. Kumar *et al.* (2011) explained that gram-positive bacteria are more resistant than gram-negative bacteria to high UV exposure. The resistance of bacteria to extreme environments is influenced by the presence of spores, pigments and high amounts of Guanine (G) and Cytosine (C), considering that bacteria with endospores and DNA with high G and C content are more resistant to temperature, humidity, light intensity, UV rays, chemicals, and free radicals.

The advantages of this research are the locations that have never been studied and the research variables are large in number. The limitation of this study is the identification of bacterial and fungal species in the Manggarai station area has not been carried out

4. CONCLUSIONS

This study concludes that there is a correlation between temperature, humidity, and light intensity in the number of bacterial colonies in the Manggarai station with a weak category. As for further researchers, besides analyzing the correlation between environmental factors and the number of bacterial colonies, it is also recommended to identify bacteria and fungi in the Manggarai station environment.

5. ACKNOWLEDGMENTS

The researchers would like to thank the Head of UPT Station Manggarai who has permitted a research site

6. REFERENCES

- Bragoszewska E, Pastuszka J. Influence of Meteorological Factors on The Level and Characteristics of Culturable Bacteria in The Air in Gliwice, Upper Silesia (Poland). Silesian, Poland: Springer US; 2018. p. 241–55. <https://doi.org/10.1007/s10453-018-9510-1>
- Cho, E.M., Hong, H.J., Park, S.H., Yoon D.K., Goung, S.J.N., Lee, C.M. Distribution and Influencing Factors of Airborne Bacteria in Public Facilities Used by Pollution-Sensitive Population: A Meta-analysis. *Int J Environ Res Public Health*. 2019;16(9). <http://dx.doi.org/10.3390/ijerph16091483>
- Liu, H., Zhang, X., Zhang, H., Yao, X., Zhou, M., He, Z., Zhang, H., Lou, L., Mao, W., Zheng, P., Hu, B. Effect Of Air Pollution On The Total Bacteria And Pathogenic Bacteria In Different Sizes Of Particulate Matter. *Environmental Pollution*. 2018;233:483–93. <https://doi.org/10.1016/j.envpol.2017.10.070>

- Kallawicha, K., Lung, S.C.C., Chuang, Y.C., Wu Da, C., Chen Hsuan, T., Tsai Y.J. Chao, J.H. Spatiotemporal Distributions And Land-Use Regression Models Of Ambient Bacteria And Endotoxins In The Greater Taipei Area. *Aerosol Air Quality Research*. 2015;15(4):1448–59.
- Jones, A.M. and Harrison, R.M. The Effects Of Meteorological Factors On Atmospheric Bioaerosol Concentrations - A Review. *Science of Total Environment*. 2004;326(1–3):151–80. doi:10.1016/j.scitotenv.2003.11.021
- Mentese, S., Arisoy, M., Rad, Y.A., Güllü, G. Bacteria and Fungi Levels in Various Indoor And Outdoor Environments In Ankara, Turkey. *Clean Journal* 2009;37(6):487–93. DOI: 10.100/c1en.2008002200
- Wu, Y.H., Chan, C.C., Chew, G.L., Shih, P.W., Lee Te, C., Chao, J.H. Meteorological Factors And Ambient Bacterial Levels in A Subtropical Urban Environment. *Int J Biometeorol*. 2012;56(6):1001–9. DOI 10.1007/s00484-011-0514-6
- Kumar, P, Mahor, P, Goel, A.K., Kamboj, D.V., Kumar O. Aero-microbiological Study On Distribution Pattern Of Bacteria and Fungi During Weekdays At Two Different Locations In Urban Atmosphere Of Gwalior, Central India. *Sci. Res. Essays*. 2011: 6(25):5435–41. DOI : 10.5897/SRE11.1485
- Goudarzi, G., Soleimani, Z., Sadeghinejad, B., Alighardashi, M., Latifi, S.M., Moradi, M. Visiting Hours Impact on Indoor to Outdoor Ratio of Fungi Concentration at Golestan University Hospital in Ahvaz, Iran. *Environ. Pollut*. 2017;6(1):62. <https://doi.org/10.5539/ep.v6n1p62>
- Saadati, Z., Shahryari, T, Ramazani, A.A., Sahlabadi, F., Nikoomanesh, F., Namaie, M.H. International Journal of Molecular and Clinical Microbiology Investigation of Indoor and Outdoor Fungal Bioaerosols and Environmental Factors in Indoor Air Quality of Nursery Schools. *Int. J. Mol. Clin. Microbiol*. 2022;12(1):1596–604.
- Zhong, X., Qi, J., Li, H., Dong, L. and Gao, D. Seasonal Distribution Of Microbial Activity In Bioaerosols In The Outdoor Environment Of The Qingdao Coastal Region. *Atmos. Environ*. 2016;140:506–13. <http://dx.doi.org/10.1016/j.atmosenv.2016.06.034>.
- Hwang, S. H. and Yoon, C.S. Main Environmental Factors Affecting Concentrations Of Culturable Airborne Bacteria in Indoor Laboratories Over A Period of One Year. *Appl. Ecol. Environ. Res*. 2017;15(1):321–33. http://dx.doi.org/10.15666/aeer/1501_321333
- Hiwar, W, King, M.F., Shuweihdi, F., Fletcher, L.A., Dancer, S.J., Noakes, C.J. What is The Relationship Between Indoor Air Quality Parameters and Airborne Microorganisms in Hospital Environments? A Systematic Review And Meta-Analysis. *Indoor Air*. 2021.31(5):1308–22. DOI: 10.1111/ina.12846
- Kamel, H.F, Saeed, H.C. and Hassan Ismael, N. Comparative Effect Of Different Visible Light Energy On Bacterial Growth. *Int. J. Adv. Res*. 2016;4(2):263–70. <http://www.journalijar.com>.
- Fithri, N. K., Handayani, P. dan Vionalita, G. Faktor-Faktor yang Berhubungan dengan Jumlah Mikroorganisme Udara Dalam Ruang Kelas Lantai 8 Universitas Esa Unggul. *Forum Ilm*. 2016;13(1):15–6.
- Gola, M., Settimo, G. and Capolongo, S. Indoor Air Quality in Inpatient Environments: A Systematic Review on Factors that Influence Chemical Pollution in Inpatient Wards. *J. Health. Eng*. 2019;2019. <https://doi.org/10.1155/2019/8358306>
- Gladka, G., Hovorukha, V., Romanovskaya, V., Tashyrev, O. Correlation Between Resistance to UV Irradiation and The Taxonomic Position of Microorganisms. *Environ. Res. Eng. Manag*. 2021 : 77(1):67–75. DOI 10.5755/j01.erem.77.1.23832
- Santos, A. L, Lopes, S., Baptista, I., Henriques, I., Gomes, N.C.M., Almeida A., Correia, A. and Cunha, A. Diversity in UV Sensitivity and recovery Potential Among Bacterioneuston And Bacterioplankton Isolates. *Lett Appl Microbiol*. 2011;52(4):360–6. doi:10.1111/j.1472-765X.2011.03011.x
- Mohana, D., Thippeswamy, S. and Abhishek, R.. Antioxidant, antibacterial, and Ultraviolet-Protective Properties of carotenoids Isolated From *Micrococcus* sp. *Radiat. Prot. Environ*. 2013;36(4):168.
- Fang, Z., Ouyang, Z., Zheng, H., Wang, X., Hu, L. Culturable Airborne Bacteria In Outdoor Environments in Beijing, China. *Microb. Ecol*. 2007;54(3):487–96. DOI: 10.1007/s00248-007-9216-3
- Maier, R.M., Palmer, M.W., Andersen, G.L., Marilyn, J., Halonen, Karen, C., Josephson, Robert, S. Maier, Fernando, D. Martinez, Julia, W. Neilson, Debra, A. Stern, Donata Vercelli and Anne, L. Wright. Environmental Determinants Of And Impact On Childhood Asthma By The Bacterial Community In Household Dust. *Appl. Environ. Microbiol*. 2010;76(8):2663–7. doi:10.1128/AEM.01665-09